arXiv:math/0510332v1 [math.GM] 16 Oct 2005

,

THERE IS AT LEAST ONE PAIR OF DOUBLE PRIMES FOR ANY EVEN NUMBER

SHOUYU DU AND ZHANLE DU

This paper is dedicated to Prof. Wei WANG.

Abstract. We proved that any even number not less than 6 can be expressed as the sum of two old primes, $2n = p_i + p_j$.

1. Introduction

The Goldbach conjecture was one of the oldest unsolved problems in number theory[3, 4]. It states that for any even number $2n$ there exists a pair of double primes (p_i, p_j) such that $2n = p_i + p_j$, and usually represented as '1+1'. The best result is the Chen's Theorem [1, 2] that every 'large' even number may be written as the sum of a prime and a semiprime (2-almost primes). Our result is as theorem 1.1.

Theorem 1.1. *There is at least one pair of double primes for any even number* $2n = p_i + p_j \geq 6$.

Let P=$\{p_1, p_2, ..., p_v\} = \{2, 3, ..., p_v\}$ be the primes not exceeding $\sqrt{2n}$, then the number of primes not exceeding $2n$ [5] is,

$$\pi(2n) = \left\{ \begin{array}{l} (\pi(\sqrt{2n}) - 1) + 2n - \left(\left\lfloor \frac{2n}{p_1} \right\rfloor + \left\lfloor \frac{2n}{p_2} \right\rfloor + \cdots + \left\lfloor \frac{2n}{p_v} \right\rfloor \right) \\ + \left(\left\lfloor \frac{2n}{p_1 p_2} \right\rfloor + \left\lfloor \frac{2n}{p_1 p_3} \right\rfloor + \cdots + \left\lfloor \frac{2n}{p_{v-1} p_v} \right\rfloor \right) - \cdots , \end{array} \right. \tag{1.1}$$

For simplicity, we can write it as,

$$\begin{aligned} \pi(2n) &= (\pi(\sqrt{2n}) - 1) + 2n \left[1 - \frac{1}{p_1}\right] \left[1 - \frac{1}{p_2}\right] \cdots \left[1 - \frac{1}{p_v}\right] \\ &= (\pi(\sqrt{2n}) - 1) + 2n \prod_{i=1}^{v} \left[1 - \frac{1}{p_i}\right], \end{aligned} \tag{1.2}$$

where $2n \left[1 - \frac{1}{p_i}\right] = 2n - \left\lfloor \frac{2n}{p_i} \right\rfloor$, $2n \left[1 - \frac{1}{p_i}\right] \left[1 - \frac{1}{p_j}\right] = 2n - \left\lfloor \frac{2n}{p_i} \right\rfloor - \left\lfloor \frac{2n}{p_j} \right\rfloor + \left\lfloor \frac{2n}{p_i p_j} \right\rfloor$. The operator $\left[1 - \frac{1}{p_i}\right]$ will leave the items which are not multiples of p_i.

In this paper, $\lfloor x \rfloor \leq x$ is the floor function, $\lceil x \rceil \geq x$ the ceiling function of x. The integral operator '[ ]', which is not the floor function in this paper, has meanings only operating on (real) number: $m[\,] = [m] = \lfloor m \rfloor$.

Date: Oct.3, 2005.

2000 *Mathematics Subject Classification.* 11A41; 11P32; 11N36 .

Key words and phrases. prime, floor function, ceiling function, integral operator, Goldbach's conjecture.

All rights reserved. This is one of a serial works, and was supported financially by Prof. Wei WANG and on loan in part.

Corresponding author: Zhanle Du.

2. The number of double prime pairs in 2n

Let $Z = \{1, 2, ..., m\} (m \leq 2n-1)$ be a natural arithmetic progression, $Z' = 2n - Z = \{2n-1, 2n-2, ..., 2n-m\}$ be its accompanying arithmetic progression, so that $2n = Z_k + Z'_k, k = 1, 2, \cdots, m$. There are m such pairs.

$$\left\{ \begin{array}{lllllll} Z & = \{ & 1, & 2, & \cdots, & m & \} \\ Z' & = \{ & 2n-1, & 2n-2, & \cdots, & 2n-m & \}. \end{array} \right. \tag{2.1}$$

After deleted all the pairs in which one or both items Z_k and Z'_k are multiples of the primes $p_i \leq \sqrt{2n}$, then the pairs left are all prime pairs or [1,2n-1].

For a certain p_i, we first delete the multiples of p_i in set Z, or the items of $Z_k \mathbf{mod} p_i = 0$,

$$y(p_i) \equiv m \left[\frac{1}{p_i} \right] = \left\lfloor \frac{m}{p_i} \right\rfloor . \tag{2.2}$$

Note the operator $\left[\frac{1}{p_i} \right]$ has meaning only operated on integer (m).

If

$$\lambda_i = (2n) \mathbf{mod} p_i \tag{2.3}$$

is not zero, we should delete the multiples of p_i in Z', i.e. the items of $(2n - Z_k) \mathbf{mod} p_i = 0$ or $Z_k \mathbf{mod} p_i = (2n) \mathbf{mod} p_i$ in set Z,

$$y'(p_i) \equiv m \left[\frac{\widetilde{1}}{p_i} \right] = \left\lfloor \frac{m + (p_i - 1) - (2n-1) \mathbf{mod} p_i}{p_i} \right\rfloor , \tag{2.4}$$

where $0 \leq (2n-1) \mathbf{mod} p_i \leq (p_i - 1)$ is the remainder modulo p_i. $0 \leq \theta_i \overset{\text{def}}{=} (p_i - 1) - (2n-1) \mathbf{mod} p_i = (p_i - \lambda_i) \leq (p_i - 1)$ is the number of elements which, when added before Z', will make a new progression $(Z'' = \{Z_a, Z'\}, Z_a = \{p_i - 1, p_i - 2, ..., \lambda_i\})$ having items of $Z''_k \mathbf{mod} p_i = 0$ in the positions of $k \mathbf{mod} p_i = 0$. If $\lambda_i \neq 0$, then it is the position of the first item with $Z'_k \mathbf{mod} p_i = 0$, and then $\theta_i = p_i - \lambda_i$. Eq. (2.2) will delete all pairs with $Z_k \mathbf{mod} p_i = 0$ in set Z, and Eq. (2.4) will delete all pairs with $Z_k \mathbf{mod} p_i = \lambda_i$ in set Z.

If $(2n) \mathbf{mod} p_i = 0$, then for some $Z_k \mathbf{mod} p_i = 0$, we have $Z'_k \mathbf{mod} p_i = (2n - Z_k) \mathbf{mod} p_i = 0$. These two items are in the same pair and should be deleted only once, thus,

$$y'(p_i) \quad = 0 \qquad \text{if} \qquad (2n) \mathbf{mod} p_i = 0. \tag{2.5}$$

When $(2n) \mathbf{mod} p_i \neq 0$, $1 \leq \theta_i \leq p_i - 1$,

$$y'(p_i) = m \left[\frac{\widetilde{1}}{p_i} \right] = m \left\lfloor \frac{1}{p_i} \right\rfloor + \delta, \tag{2.6}$$

where $0 \leq \delta \leq 1$ with,

$$\delta = \left\{ \begin{array}{ll} 0: & 0 \leq m \mathbf{mod} p_i + \theta_i < p_i \\ 1: & \text{else.} \end{array} \right. \tag{2.7}$$

After deleted all the multiples of p_i in both Z and Z', the pairs will leave,

$$M(p_i) = m - y(p_i) - y'(p_i) \equiv m \left[1 - \frac{1}{p_i} - \frac{\widetilde{1}}{p_i} \right] . \tag{2.8}$$

The operator $\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right]$, when operating on m, will leaves the items having no multiples of p_i. After deleted the multiples of the primes $p_i \le p_v$ in both Z and Z', the pairs left will be prime pairs and have,

$$\begin{aligned} D_0(m) &= m\left[1-\frac{1}{p_1}\right]\left[1-\frac{1}{p_2}-\frac{\widetilde{1}}{p_2}\right]\cdots\left[1-\frac{1}{p_v}-\frac{\widetilde{1}}{p_v}\right] \\ &= m\prod_{(2n)\mathbf{mod}p_i=0}\left[1-\frac{1}{p_i}\right]\prod_{(2n)\mathbf{mod}p_j\neq0}\left[1-\frac{1}{p_j}-\frac{\widetilde{1}}{p_j}\right], \end{aligned} \tag{2.9}$$

the meaning is as follows,

$$\begin{aligned} & m\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right]\left[1-\frac{1}{p_j}-\frac{\widetilde{1}}{p_j}\right] \\ = \ & m\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}-\frac{1}{p_j}+\frac{1}{p_ip_j}+\frac{\widetilde{1}}{p_i}\frac{1}{p_j}-\frac{\widetilde{1}}{p_j}+\frac{1}{p_i}\frac{\widetilde{1}}{p_j}+\frac{\widetilde{1}}{p_i}\frac{\widetilde{1}}{p_j}\right], \end{aligned} \tag{2.10}$$

$$\begin{cases} m\left[\frac{1}{p_i}\right] = \left\lfloor\frac{m}{p_i}\right\rfloor \\ m\left[\frac{\widetilde{1}}{p_i}\right] = \left\lfloor\frac{m+(p_i-1)-(2n-1)\mathbf{mod}p_i}{p_i}\right\rfloor = \left\lfloor\frac{m+\theta_i}{p_i}\right\rfloor \left\lfloor\frac{m+p_i-\lambda_i}{p_i}\right\rfloor \\ m\left[\frac{1}{p_i}\right]\left[\frac{1}{p_j}\right] = m\left[\frac{1}{p_ip_j}\right] = \left\lfloor\frac{m}{p_ip_j}\right\rfloor \\ m\left[\frac{\widetilde{1}}{p_i}\right]\left[\frac{\widetilde{1}}{p_j}\right] = m\left[\frac{\widetilde{1}}{p_ip_j}\right] = \left\lfloor\frac{m+(p_ip_j-1)-(2n-1)\mathbf{mod}p_ip_j}{p_ip_j}\right\rfloor = \left\lfloor\frac{m+p_ip_j-\lambda_{ij}}{p_ip_j}\right\rfloor \\ m\left[\frac{1}{p_i}\right]\left[\frac{\widetilde{1}}{p_j}\right] = m\left[\frac{1}{p_i}\frac{\widetilde{1}}{p_j}\right] = \left\lfloor\frac{m+\theta_{i,j}}{p_ip_j}\right\rfloor = \left\lfloor\frac{m+p_ip_j-\lambda_{i,j}}{p_ip_j}\right\rfloor = \left\lfloor\frac{\left\lfloor\frac{m}{p_i}\right\rfloor+p_j-\lambda_{i|j}}{p_j}\right\rfloor. \end{cases} \tag{2.11}$$

Let $1 \le \theta_{ij} \overset{\text{def}}{=} p_ip_j - 1 - (2n-1)\mathbf{mod}p_ip_j = p_ip_j - \lambda_{ij} \le (p_ip_j - 1)$. $1 \le \lambda_{ij} \overset{\text{def}}{=} (2n)\mathbf{mod}p_ip_j \le p_ip_j - 1$ is the position of the first item in Z' with $Z'_k\mathbf{mod}(p_ip_j) = 0$, if it does not exist then $\lambda_{ij} > m, m + \theta_{ij} < p_ip_j$ and this item will be zero. Note that $\lambda_{ij} \neq 0$ because $p_i \nmid (2n)$ and $p_j \nmid (2n)$.

In Eq. (2.11), $1 \le \theta_{i,j} \overset{\text{def}}{=} p_ip_j - \lambda_{i,j} \le (p_ip_j - 1)$, $1 \le \lambda_{i,j} \le p_ip_j - 1$ is the position of the first item with $p_i | Z_{\lambda_{i,j}}$ and $p_j | Z'_{\lambda_{i,j}}$.

$$\begin{cases} Z_{\lambda_{i,j}}\mathbf{mod}p_i = 0 \\ Z_{\lambda_{i,j}}\mathbf{mod}p_j = (2n)\mathbf{mod}p_j. \end{cases} \tag{2.12}$$

Let $X = \{\lambda p_i\}$, $X' = 2n - X = \{2n - \lambda p_i\}$, $\lambda = 1, 2, \cdots, \left\lfloor\frac{m}{p_i}\right\rfloor$, then

$$\begin{cases} \lambda_{i|j}p_i\mathbf{mod}p_j &= (2n)\mathbf{mod}p_j \\ \lambda_{i,j} &= \lambda_{i|j}p_i. \end{cases} \tag{2.13}$$

If it exists and $(2n)\mathbf{mod}p_j \neq 0$, then $1 \le \lambda_{i|j} \le p_j - 1$, and $p_i \le \theta_{i,j} = p_ip_j - \lambda_{i,j} \le p_i(p_j - 1)$. If there is no such $\lambda_{i,j}$ in set Z, i.e., $\lambda_{i,j} \ge (m+1)$, then the last item in Eq. (2.11) will equal zero.

Let $m = 2n-1$, then $Z = \{1, 2, ..., 2n-1\}$ and $Z' = 2n - Z = \{2n-1, 2n-2, ..., 1\}$. For each pais, there is another same pair: $2n = Z_k + Z'_k = Z'_{2n-k} + Z_{2n-k}$. When $k = n$ is prime, it must be the same pair $2n = Z_n + Z'_n = n + n$. Therefore, from

Eq. (2.9), the actual number of prime pairs in 2n is,

$$
\begin{aligned}
D(2n) &= \left\lceil \tfrac{D_0(2n-1)}{2} \right\rceil + D(\sqrt{2n}) - D_1 \\
&= \left\lceil \tfrac{1}{2}(2n-1)\left[1-\tfrac{1}{2}\right]\prod_{i=2}^{v}\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right]\right\rceil + D(\sqrt{2n}) - D_1,
\end{aligned} \tag{2.14}
$$

where $D(\sqrt{2n}) \geq 0$ is the number of prime pairs $2n = p_i + (2n - p_i)$ when $p_i \leq p_v < \sqrt{2n}$, $p_j = 2n - p_i$ are both primes, and

$$
D_1 = \begin{cases} 0 & 2n-1 \neq prime \\ 1 & 2n-1 = prime. \end{cases} \tag{2.15}
$$

Remember that when $(2n)\mathbf{mod}p_i = 0$, $\left[\frac{\widetilde{1}}{p_i}\right] = 0$. □

Example 2.1. Let $2n = 46$, then $p_i = [2, 3, 5]$, $3 \nmid n, 5 \nmid n$, $m = 2n - 1 = 45$,

$$
\begin{cases} Z &= \{ \quad 1 \quad 2 \quad 3 \quad \cdots \quad 23 \quad \cdots \quad 44 \quad 45\} \\ Z' &= \{ \quad 45 \quad 44 \quad 43 \quad \cdots \quad 23 \quad \cdots \quad 2 \quad 1\} \end{cases} \tag{2.16}
$$

From Eq. (2.3),

$$
\begin{cases} p_i & : \quad 3 \quad 5 \quad 15 \\ \lambda_i & : \quad 1 \quad 1 \quad 1. \end{cases} \tag{2.17}
$$

From Eq. (2.13),

$$
\begin{cases} (p_i, p_j) & : \quad (2,3) \quad (5,3) \quad (10,3) \quad (2,5) \quad (3,5) \quad (6,5) \quad (2,15) \\ \lambda_{i,j} & : \quad 4 \quad 10 \quad 10 \quad 6 \quad 6 \quad 6 \quad 16) \end{cases} \tag{2.18}
$$

From Eq. (2.9), (2.11),

$$
\begin{aligned}
D_0(2n-1) &= (2n-1)\left[1-\tfrac{1}{2}\right]\left[1-\tfrac{1}{3}-\tfrac{\widetilde{1}}{3}\right]\left[1-\tfrac{1}{5}-\tfrac{\widetilde{1}}{5}\right] \\
&= 45\Big[1-\tfrac{1}{2}-\tfrac{1}{3}+\tfrac{1}{6}-\tfrac{\widetilde{1}}{3}+\tfrac{1}{2}\tfrac{\widetilde{1}}{3}-\tfrac{1}{5}+\tfrac{1}{10}+\tfrac{1}{15}-\tfrac{1}{30}+\tfrac{1}{5}\tfrac{\widetilde{1}}{3}-\tfrac{1}{10}\tfrac{\widetilde{1}}{3} \\
&\quad -\tfrac{\widetilde{1}}{5}+\tfrac{1}{2}\tfrac{\widetilde{1}}{5}+\tfrac{1}{3}\tfrac{\widetilde{1}}{5}-\tfrac{1}{6}\tfrac{\widetilde{1}}{5}+\tfrac{\widetilde{1}}{15}-\tfrac{1}{2}\tfrac{\widetilde{1}}{15}\Big] \\
&= 45-\left\lfloor\tfrac{45}{2}\right\rfloor-\left\lfloor\tfrac{45}{3}\right\rfloor+\left\lfloor\tfrac{45}{6}\right\rfloor-\left\lfloor\tfrac{45+3-1}{3}\right\rfloor+\left\lfloor\tfrac{45+6-4}{6}\right\rfloor \\
&\quad -\left\lfloor\tfrac{45}{5}\right\rfloor+\left\lfloor\tfrac{45}{10}\right\rfloor+\left\lfloor\tfrac{45}{15}\right\rfloor-\left\lfloor\tfrac{45}{30}\right\rfloor+\left\lfloor\tfrac{45+15-10}{15}\right\rfloor-\left\lfloor\tfrac{45+30-10}{30}\right\rfloor \\
&\quad -\left\lfloor\tfrac{45+5-1}{5}\right\rfloor+\left\lfloor\tfrac{45+10-6}{10}\right\rfloor+\left\lfloor\tfrac{45+15-6}{15}\right\rfloor-\left\lfloor\tfrac{45+30-6}{30}\right\rfloor \\
&\quad +\left[\tfrac{45+15-1}{15}\right]-\left\lfloor\tfrac{45+30-16}{30}\right\rfloor \\
&= 45-22-15+7-15+7 \\
&\quad -9+4+3-1+3-2 \\
&\quad -9+4+3-2+3-1 \\
&= 3.
\end{aligned}
$$

We can check this result directly. After deleted the items of $Z_k\mathbf{mod}2 = 0; Z_k\mathbf{mod}3 = 0, 1; Z_k\mathbf{mod}5 = 0, 1$ from set Z, it is left $Z = \{17, 23, 29\}$. $D_0(2n-1) = 3$, it is the same as before.

From $D(\sqrt{2n}) = 2(46 = 3 + 43 = 5 + 41)$, $D_1 = 0$(46-1=45 is not a prime). From Eq. (2.14), $D(2n) = \lceil\frac{3}{2}\rceil + 2 - 0 = 4$. The set left is $Z = \{3, 5, 17, 23\}, Z' =$

$44 - Z = \{43, 41, 29, 23\}$. Thus there are four double prime pairs of $46 = p_i + p_j$: $(3, 43), (5, 41), (17, 29), (23, 23)$. □

Definition 2.2. The items of $Z_k \mathbf{mod} p_i = 0$, or the multiples of p_i, have,

$$S(m, p_i \parallel 0) =: \sum_{Z_k \mathbf{mod} p_i = 0} 1 = m \left[\tfrac{1}{p_i} \right] = \left\lfloor \tfrac{m}{p_i} \right\rfloor . \tag{2.19}$$

So the items of $Z_k \mathbf{mod} p_j \neq 0, \lambda_j$ have,

$$S(m, p_j \nparallel 0, \lambda_j) \quad =: \sum_{Z_k \mathbf{mod} p_i \neq 0, \lambda_j} 1 = m \left[1 - \tfrac{1}{p_j} - \tfrac{\tilde{1}}{p_j} \right] . \tag{2.20}$$

Definition 2.3. Let $X = \{X_1, X_2, \cdots, X_t\}$ be an (any) integer set, then

$$t \left[1 - \tfrac{1'}{p_i} - \tfrac{1''}{p_i} \right] =: \sum_{X_k \mathbf{mod} p_i \neq 0, \lambda_i} 1 = t - \left[\tfrac{t + \phi_i'}{p_i} \right] - \left[\tfrac{t + \phi_i''}{p_i} \right] \geq 0 \tag{2.21}$$

is the number of items left after deleted the items of $X_k \mathbf{mod} p_i = 0, \lambda_i$.

Usually $m \left[1 - \frac{1}{p_i} - \frac{\tilde{1}}{p_i} \right] \left[1 - \frac{1}{p_j} - \frac{\tilde{1}}{p_j} \right] \neq \left(m \left[1 - \frac{1}{p_i} - \frac{\tilde{1}}{p_i} \right] \right) \left[1 - \frac{1}{p_j} - \frac{\tilde{1}}{p_j} \right]$, We can express it as

$$\begin{aligned} S(m, p_i \nparallel 0, \lambda_i; p_j \nparallel 0, \lambda_j) &= m \left[1 - \tfrac{1}{p_i} - \tfrac{\tilde{1}}{p_i} \right] \left[1 - \tfrac{1}{p_j} - \tfrac{\tilde{1}}{p_j} \right] \\ &= \left(m \left[1 - \tfrac{1}{p_i} - \tfrac{\tilde{1}}{p_i} \right] \right) \left[1 - \tfrac{1'}{p_j} - \tfrac{1''}{p_j} \right] , \end{aligned} \tag{2.22}$$

is the number of items left when we first delete those $Z_k \mathbf{mod} p_i = 0, \lambda_i$ from set Z, and then delete those $X_{k'} \mathbf{mod} p_j = 0, \lambda_j$ from set $X = \{Z, X_{k'} \mathbf{mod} p_i \neq 0, \lambda_i, k' = 1, 2, \cdots, m \left[1 - \frac{1}{p_i} - \frac{\tilde{1}}{p_i} \right] \}$, where set X is no longer an arithmetic sequence. In general,

$$m \prod_{i=1}^{i_m} \left[1 - \tfrac{1}{p_i} - \tfrac{\tilde{1}}{p_i} \right] \left[1 - \tfrac{1}{p_j} - \tfrac{\tilde{1}}{p_j} \right] = \left(m \prod_{i=1}^{i_m} \left[1 - \tfrac{1}{p_i} - \tfrac{\tilde{1}}{p_i} \right] \right) \left[1 - \tfrac{1'}{p_j} - \tfrac{1''}{p_j} \right] . \tag{2.23}$$

3. Some property

$$\begin{cases} \left[1 - \tfrac{1}{p_i} \right] \left[1 - \tfrac{1}{p_j} \right] = \left[1 - \tfrac{1}{p_j} \right] \left[1 - \tfrac{1}{p_i} \right] \\ \left[1 - \tfrac{1}{p_i} - \tfrac{\tilde{1}}{p_i} \right] \left[1 - \tfrac{1}{p_j} - \tfrac{\tilde{1}}{p_j} \right] = \left[1 - \tfrac{1}{p_j} - \tfrac{\tilde{1}}{p_j} \right] \left[1 - \tfrac{1}{p_i} - \tfrac{\tilde{1}}{p_i} \right] . \end{cases} \tag{3.1}$$

$$\left[\tfrac{m}{p_i} \right] \quad = \left[\tfrac{m_1 + m_2}{p_i} \right] = \left[\tfrac{m_1}{p_i} \right] + \left[\tfrac{m_2}{p_i} \right] + \left[\tfrac{m_1 \mathbf{mod} p_i + m_2 \mathbf{mod} p_i}{p_i} \right] . \tag{3.2}$$

$$m \left[1 - \tfrac{1}{p_i} \right] \geq m \left[1 - \tfrac{1}{p_i} - \tfrac{\tilde{1}}{p_i} \right] \geq 0. \tag{3.3}$$

$$m \left[1 - \tfrac{1}{p_i} - \tfrac{\tilde{1}}{p_i} \right] = a p_i \left(1 - \tfrac{2}{p_i} \right) + b \left[1 - \tfrac{1}{p_i} - \tfrac{\tilde{1}}{p_i} \right] \quad \text{for } m = a p_i + b. \tag{3.4}$$

$$-2 \leq (m_1 + m_2) \left[1 - \tfrac{1}{p_i} - \tfrac{\tilde{1}}{p_i} \right] - m_1 \left[1 - \tfrac{1}{p_i} - \tfrac{\tilde{1}}{p_i} \right] - m_2 \left[1 - \tfrac{1}{p_i} - \tfrac{\tilde{1}}{p_i} \right] \leq 1. \tag{3.5}$$

Proof. Let $\alpha = m_1\mathbf{mod}p_i, \beta = m_2\mathbf{mod}p_i$,

$$\begin{array}{rcl}\delta_{12} & = & (m_1+m_2)\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right]-m_1\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right]-m_2\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right] \\ & = & m_1+m_2-\left[\tfrac{m_1+m_2}{p_i}\right]-\left[\tfrac{m_1+m_2+\theta_i}{p_i}\right]-m_1+\left[\tfrac{m_1}{p_i}\right]+\left[\tfrac{m_1+\theta_i}{p_i}\right] \\ & & -m_2+\left[\tfrac{m_2}{p_i}\right]+\left[\tfrac{m_2+\theta_i}{p_i}\right] \\ & = & -\left[\tfrac{\alpha+\beta}{p_i}\right]-\left[\tfrac{\alpha+\beta+\theta_i}{p_i}\right]+\left[\tfrac{\alpha+\theta_i}{p_i}\right]+\left[\tfrac{\beta+\theta_i}{p_i}\right].\end{array}$$

The minimum: $min(\delta_{12}) = -1-1 = -2$, when $\alpha+\beta \geq p_i$, $\alpha+\theta_i < p_i$, $\beta+\theta_i < p_i$.

The maximum: $max(\delta_{12}) = -0-1+1+1 = 1$, when $\alpha+\beta < p_i$, $\alpha+\theta_i \geq p_i$, $\beta+\theta_i \geq p_i$. □

We can represent Eq. (3.5) as

$$\begin{array}{rl} & (m_1+m_2)\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right] \\ = & m_1+m_2-\left[\tfrac{m_1+m_2}{p_i}\right]-\left[\tfrac{m_1+m_2+p_i-1-(2n-1)\mathbf{mod}p_i}{p_i}\right] \\ = & m_1-\left[\tfrac{m_1}{p_i}\right]-\left[\tfrac{m_1+p_i-1-(2n-1)\mathbf{mod}p_i}{p_i}\right] \\ & +m_2-\left[\tfrac{m_1\mathbf{mod}p_i+m_2}{p_i}\right]-\left[\tfrac{m_2+p_i-1-(2n-1-m_1)\mathbf{mod}p_i)}{p_i}\right] \\ \equiv & m_1\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right]+m_2\left[1-\tfrac{1'}{p_i}-\tfrac{1''}{p_i}\right].\end{array} \tag{3.6}$$

$m_2\left[\tfrac{1'}{p_i}\right]$ will delete the items of $X_k\mathbf{mod}p_i = 0$ in the sequence of $X = \{m_1+1, m_1+2, \cdots, m_1+m_2\}$, and $m_2\left[\tfrac{1''}{p_i}\right]$ will delete the items of $X'_k\mathbf{mod}p_i = 0$ in the sequence of $X' = \{2n-X\}$ or the items of $X_k\mathbf{mod}p_i = \lambda_i$ in set X.

For any m_2, $0 \leq m_2\left[1-\tfrac{1'}{p_i}-\tfrac{1''}{p_i}\right] \leq m_2$, so,

$$m'\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right] \geq m\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right] \quad \text{for} \quad m' \geq m. \tag{3.7}$$

4. SOME LEMMA

Lemma 4.1. *For $m \geq p_i$,*

$$m\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right] \geq \left\lceil m\left(1-\tfrac{3}{p_i}\right)\right\rceil. \tag{4.1}$$

Proof.

$$\begin{array}{rl} left = \left\lceil m-\left\lfloor\tfrac{m}{p_i}\right\rfloor-\left\lfloor\tfrac{m+\theta_i}{p_i}\right\rfloor\right\rceil & \geq \left\lceil m-\left\lfloor\tfrac{m}{p_i}\right\rfloor-\left\lfloor\tfrac{m}{p_i}\right\rfloor-1\right\rceil \\ & \geq \left\lceil m-\tfrac{2m}{p_i}-\tfrac{m}{p_i}\right\rceil = right.\end{array}$$

□

Lemma 4.2. *For $m \geq p_j^2$, $p_j > p_i \geq 2$,*

$$m\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right]\left[1-\tfrac{1}{p_j}-\tfrac{\widetilde{1}}{p_j}\right] \geq \left\lceil m\left(1-\tfrac{3}{p_j}\right)\right\rceil\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right]. \tag{4.2}$$

Proof. Let $m = sp_ip_j+t$, $t = ap_j+b$, $0 \leq a \leq (p_i-1),\ 0 \leq b \leq (p_j-1)$, Suppose $\lambda_i = (2n)\mathbf{mod}p_i \neq 0$ else $\left[\tfrac{\widetilde{1}}{p_i}\right] = 0$, and $\lambda_j = (2n)\mathbf{mod}p_j \neq 0$ else $\left[\tfrac{\widetilde{1}}{p_j}\right] = 0$, so $\theta_i, \theta_j > 0$ and $\theta_{i,j}, \theta_{j,i} > 0, \theta_{ij} > 0$. Because $m \geq p_j^2,\ p_j > p_i$, so $s \geq 1$.

Because $m(v) = m \geq p_v^2$. Let $m(j-1) = \lceil m(j)(1-3/p_j) \rceil$, then

$$\begin{aligned} m(j-1) &= \lceil m(j)(1-3/p_j) \rceil \geq p_j^2(1-3/p_j) = p_j(p_j-3) \\ &\geq (p_{j-1}+2)(p_{j-1}-1) = p_{j-1}^2 + p_{j-1} - 2 \geq p_{j-1}^2. \end{aligned}$$

So for any $i \leq (j-1)$, we have $m(i) \geq p_i^2$.

From Eq. (3.2), (3.4),

$$\begin{aligned} \varepsilon \quad &= m\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right]\left[1-\tfrac{1}{p_j}-\tfrac{\widetilde{1}}{p_j}\right] - \left\lceil m\left(1-\tfrac{3}{p_j}\right)\right\rceil \left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right] \\ &= sp_ip_j\left(1-\tfrac{2}{p_i}\right)\left(1-\tfrac{2}{p_j}\right) - sp_ip_j\left(1-\tfrac{2}{p_i}\right)\left(1-\tfrac{3}{p_j}\right) \\ &\quad +t - \left\lfloor \tfrac{t}{p_i} \right\rfloor - \left\lfloor \tfrac{t+\theta_i}{p_i} \right\rfloor - \left\lfloor \tfrac{t}{p_j} \right\rfloor + \left\lfloor \tfrac{t}{p_ip_j} \right\rfloor + \left\lfloor \tfrac{t+\theta_{j,i}}{p_ip_j} \right\rfloor - \left\lfloor \tfrac{t+\theta_j}{p_j} \right\rfloor + \left\lfloor \tfrac{t+\theta_{i,j}}{p_ip_j} \right\rfloor \\ &\quad + \left\lfloor \tfrac{t+\theta_{ij}}{p_ip_j} \right\rfloor - \left(t - \left\lfloor \tfrac{3t}{p_j} \right\rfloor\right) + \left\lfloor \frac{t-\left\lfloor \frac{3t}{p_j} \right\rfloor}{p_i} \right\rfloor + \left\lfloor \frac{t+\theta_i-\left\lfloor \frac{3t}{p_j} \right\rfloor}{p_i} \right\rfloor \\ &= s(p_i-2) \\ &\quad + \left\lfloor \tfrac{3t}{p_j} \right\rfloor - \left\lfloor \tfrac{t}{p_j} \right\rfloor - \left\lfloor \tfrac{t+\theta_j}{p_j} \right\rfloor + \left\lfloor \tfrac{t}{p_ip_j} \right\rfloor + \left\lfloor \tfrac{t+\theta_{j,i}}{p_ip_j} \right\rfloor + \left\lfloor \tfrac{t+\theta_{i,j}}{p_ip_j} \right\rfloor \\ &\quad + \left\lfloor \tfrac{t+\theta_{ij}}{p_ip_j} \right\rfloor - \left\lfloor \frac{\left\lfloor \frac{3t}{p_j} \right\rfloor}{p_i} \right\rfloor - \varepsilon_1 - \left\lfloor \frac{\left\lfloor \frac{3t}{p_j} \right\rfloor}{p_i} \right\rfloor - \varepsilon_2 \\ &= s(p_i-2) \\ &\quad +3a + \left\lfloor \tfrac{3b}{p_j} \right\rfloor - a - a - \left\lfloor \tfrac{b+\theta_j}{p_j} \right\rfloor + 0 + \left\lfloor \tfrac{ap_j+b+\theta_{j,i}}{p_ip_j} \right\rfloor + \left\lfloor \tfrac{ap_j+b+\theta_{i,j}}{p_ip_j} \right\rfloor \\ &\quad + \left\lfloor \tfrac{ap_j+b+\theta_{ij}}{p_ip_j} \right\rfloor - 2\left\lfloor \frac{3a+\left\lfloor \frac{3b}{p_j} \right\rfloor}{p_i} \right\rfloor - \varepsilon_1 - \varepsilon_2 \\ &= s(p_i-2) + a + \left\lfloor \tfrac{3b}{p_j} \right\rfloor + \left\lfloor \tfrac{ap_j+b+\theta_{j,i}}{p_ip_j} \right\rfloor + \left\lfloor \tfrac{ap_j+b+\theta_{i,j}}{p_ip_j} \right\rfloor + \left\lfloor \tfrac{ap_j+b+\theta_{ij}}{p_ip_j} \right\rfloor \\ &\quad -\varepsilon_1 - \varepsilon_2 - 2\varepsilon_3 - \varepsilon_4, \end{aligned}$$

where

$$\begin{cases} \varepsilon_1 &= \left\lfloor \dfrac{\left\lfloor \frac{3t}{p_j} \right\rfloor \mathbf{mod} p_i + \left(t - \left\lfloor \frac{3t}{p_j} \right\rfloor\right) \mathbf{mod} p_i}{p_i} \right\rfloor \leq 1 \\ \varepsilon_2 &= \left\lfloor \dfrac{\left\lfloor \frac{3t}{p_j} \right\rfloor \mathbf{mod} p_i + \left(t+\theta_i - \left\lfloor \frac{3t}{p_j} \right\rfloor\right) \mathbf{mod} p_i}{p_i} \right\rfloor \leq 1 \\ \varepsilon_3 &= \left\lfloor \dfrac{3a+\left\lfloor \frac{3b}{p_j} \right\rfloor}{p_i} \right\rfloor \leq 2 \\ \varepsilon_4 &= \left\lfloor \dfrac{b+\theta_j}{p_j} \right\rfloor \leq 1, \end{cases}$$

$$\Delta\varepsilon = \varepsilon_1 + \varepsilon_2 + 2\varepsilon_3 + \varepsilon_4 \leq 7.$$

(1) If $p_i \geq 11$, then $\varepsilon \geq s(11-2) - \Delta\varepsilon \geq 9 - 7 > 0$.

(2) If $p_i = 7$, then
  (a) if $a \geq 2$, then $\varepsilon \geq s(p_i-2) + a - \Delta\varepsilon \geq 5 + 2 - 7 = 0$.
  (b) if $a \leq 1$, then $\varepsilon_3 = \left\lfloor \dfrac{3a+\left\lfloor \frac{3b}{p_j} \right\rfloor}{p_i} \right\rfloor \leq \left\lfloor \frac{3+2}{7} \right\rfloor = 0$, $\Delta\varepsilon \leq 3$, so $\varepsilon \geq s(p_i - 2) - \Delta\varepsilon \geq 5 - 3 > 0$.

(3) If $p_i = 5$,
  (a) if $a \geq 4$: $\varepsilon \geq s(p_i-2) + a - \Delta\varepsilon \geq 3 + 4 - 7 = 0$.
  (b) if $a = 0$: $\varepsilon_3 = 0$, $\Delta\varepsilon \leq 3$, so $\varepsilon \geq s(p_i-2) - \Delta\varepsilon \geq 3 - 3 = 0$.

(c) if $a = 1$: if $\left\lfloor \frac{3b}{p_j} \right\rfloor \geq 1$ then $\varepsilon_3 \leq 1$, $\Delta\varepsilon \leq 5$, so $\varepsilon \geq s(p_i - 2) + a + \left\lfloor \frac{3b}{p_j} \right\rfloor - \Delta\varepsilon \geq 3 + 1 + 1 - 5 = 0$; else for $\left\lfloor \frac{3b}{p_j} \right\rfloor = 0$ then $\varepsilon_3 = 0$, $\Delta\varepsilon \leq 3$, so $\varepsilon \geq s(p_i - 2) + a - \Delta\varepsilon \geq 3 + 1 - 3 > 0$.

(d) if $a = 2$: $\varepsilon_3 = 1$, $\Delta\varepsilon \leq 5$, so $\varepsilon \geq s(p_i - 2) + a - \Delta\varepsilon \geq 3 + 2 - 5 = 0$.

(e) if $a = 3$: if $\left\lfloor \frac{3b}{p_j} \right\rfloor \geq 1$ then $\varepsilon \geq s(p_i - 2) + a + \left\lfloor \frac{3b}{p_j} \right\rfloor - \Delta\varepsilon \geq 3 + 3 + 1 - 7 = 0$; else for $\left\lfloor \frac{3b}{p_j} \right\rfloor = 0$ then $\varepsilon_3 = \left\lfloor \frac{3a + \left\lfloor \frac{3b}{p_j} \right\rfloor}{p_i} \right\rfloor = 1$, $\Delta\varepsilon \leq 5$, so $\varepsilon \geq s(p_i - 2) + a - \Delta\varepsilon \geq 3 + 3 - 5 > 0$.

(4) If $p_i = 3$, then $0 \leq a \leq 2$,

$$\begin{cases} a - 2\varepsilon_3 & = a - 2\left\lfloor \frac{3a + \left\lfloor \frac{3b}{p_j} \right\rfloor}{3} \right\rfloor = a - 2a = -a \\ \varepsilon_1 & = \left\lfloor \frac{\left(\left\lfloor \frac{3b}{p_j} \right\rfloor\right)\mathbf{mod}3 + \left(ap_j + b - \left\lfloor \frac{3b}{p_j} \right\rfloor\right)\mathbf{mod}3}{3} \right\rfloor \\ \varepsilon_2 & = \left\lfloor \frac{\left(\left\lfloor \frac{3b}{p_j} \right\rfloor\right)\mathbf{mod}3 + \left(ap_j + b + \theta_i - \left\lfloor \frac{3b}{p_j} \right\rfloor\right)\mathbf{mod}3}{3} \right\rfloor, \end{cases}$$

$$\begin{aligned} \varepsilon &= s - a + \left\lfloor \frac{3b}{p_j} \right\rfloor + \left\lfloor \frac{ap_j + b + \theta_{j,i}}{3p_j} \right\rfloor + \left\lfloor \frac{ap_j + b + \theta_{i,j}}{3p_j} \right\rfloor + \left\lfloor \frac{ap_j + b + \theta_{ij}}{3p_j} \right\rfloor \\ &\quad - \varepsilon_1 - \varepsilon_2 - \varepsilon_4. \end{aligned}$$

(a) if $\left\lfloor \frac{3b}{p_j} \right\rfloor = 0$, then $\varepsilon_1 = \varepsilon_2 = 0$.

- $a = 0$: $\varepsilon \geq s - \varepsilon_4 \geq 0$.
- if $a \geq 1$: $\varepsilon' \equiv s + \left\lfloor \frac{ap_j + b + \theta_{i,j}}{3p_j} \right\rfloor - \varepsilon_4 \geq 1$.

Proof. – If $s \geq 2$ then $\varepsilon' \geq 1$.

– If $s = 1$ i.e., $s = \left\lfloor \frac{m}{p_i p_j} \right\rfloor \geq \left\lfloor \frac{p_j^2}{3p_j} \right\rfloor = 1$, so $p_j = 5$. From $m \geq 5^2 = 3 \cdot 5 + 2 \cdot 5$, we have $a = 2$. From $\left\lfloor \frac{3b}{p_j} \right\rfloor = 0$, we have $b \leq 1$. Because $\theta_j \leq p_j - 1$, so $b + \theta_j \leq p_j$.
If $b + \theta_j \leq p_j - 1$, i.e., $b = 0$ or $\theta_j \leq p_j - 2$, then $\varepsilon_4 = 0$.
Else for $b + \theta_j = p_j$, i.e., $b = 1$ and $\theta_j = p_j - 1$, $\varepsilon_4 = 1$. But $\theta_j = p_j - 1 - (2n-1)\mathbf{mod}p_j$, so $(2n-1)\mathbf{mod}p_j = 0$, or $(2n)\mathbf{mod}p_j = 1$.
Let us consider $\theta_{i,j} = p_i p_j - \lambda_{i,j}$. From Eq. (2.13), $\lambda_{i,j} = \lambda p_i = 3\lambda$, with the condition $(2n - 3\lambda)\mathbf{mod}p_j = 0$.
$(3\lambda)\mathbf{mod}p_j = (2n)\mathbf{mod}5 = 1$, so $\lambda = 2$ and $\theta_{i,j} = p_i p_j - 3\lambda = 3 \cdot 5 - 3 \cdot 2 = 9$. $\left\lfloor \frac{ap_j + b + \theta_{i,j}}{3p_j} \right\rfloor \geq \left\lfloor \frac{2 \times 5 + 1 + 9}{3 \cdot 5} \right\rfloor = 1$.
Therefore, $\left\lfloor \frac{ap_j + b + \theta_{i,j}}{3p_j} \right\rfloor - \varepsilon_4 \geq 0 \quad \text{if} \quad s = 1$

or $\varepsilon' = s + \left\lfloor \frac{ap_j + b + \theta_{i,j}}{3p_j} \right\rfloor - \varepsilon_4 \geq 1$.

□

$$\varepsilon = \varepsilon' - a + \left\lfloor \frac{ap_j + b + \theta_{j,i}}{3p_j} \right\rfloor + \left\lfloor \frac{ap_j + b + \theta_{ij}}{3p_j} \right\rfloor$$

If $a = 1$, then $\varepsilon \geq 1 - a \geq 0$.
Else if $a = 2(< p_i)$, from Eq. (2.13), $\theta_{j,i} = p_i p_j - \lambda p_j \geq p_j$ $(1 \leq \lambda \leq p_i - 1)$, $\left\lfloor \frac{ap_j+b+\theta_{j,i}}{p_i p_j} \right\rfloor \geq \left\lfloor \frac{2p_j+p_j}{3p_j} \right\rfloor = 1$. So $\varepsilon \geq 1 - a + 1 \geq 0$.

(b) if $\left\lfloor \frac{3b}{p_j} \right\rfloor \geq 1$, then $\varepsilon'' \equiv \left\lfloor \frac{3b}{p_j} \right\rfloor - \varepsilon_1 - \varepsilon_2 \geq 0$.

Proof.
- if $\left\lfloor \frac{3b}{p_j} \right\rfloor = 2$, then $\varepsilon'' \geq 0$.
- if $\left\lfloor \frac{3b}{p_j} \right\rfloor = 1$, because $\theta_i \geq 1$,

$$\begin{aligned}
\varepsilon_1 + \varepsilon_2 &= \left\lfloor \tfrac{1+(ap_j+b-1)\mathbf{mod}3}{3} \right\rfloor + \left\lfloor \tfrac{1+(ap_j+b+\theta_i-1)\mathbf{mod}3}{3} \right\rfloor \\
&\leq \begin{cases} 1+0=1 & \text{if} \quad (ap_j+b)\mathbf{mod}3 = 0 \\ 0+1=1 & \text{if} \quad (ap_j+b)\mathbf{mod}3 = 1 \\ 0+1=1 & \text{if} \quad (ap_j+b)\mathbf{mod}3 = 2. \end{cases}
\end{aligned}$$

So $\varepsilon_1 + \varepsilon_2 \leq 1$ and $\varepsilon'' = \left\lfloor \frac{3b}{p_j} \right\rfloor - \varepsilon_1 - \varepsilon_2 \geq 0$.

□

Besides, $\varepsilon' \equiv s + \left\lfloor \frac{ap_j+b+\theta_{i,j}}{3p_j} \right\rfloor - \varepsilon_4 \geq 1$.

Proof.
- If $s \geq 2$ then $\varepsilon' \geq 1$.
- If $s = 1$ i.e., $s = \left\lfloor \frac{m}{p_i p_j} \right\rfloor \geq \left\lfloor \frac{p_j^2}{3p_j} \right\rfloor = 1$, so $p_j = 5$. From $5^2 = 3 \cdot 5 + 2 \cdot 5$, we have $a = 2$. From $\left\lfloor \frac{3b}{p_j} \right\rfloor = 1, 2$, we have $b = 2, 3, 4$. Let us consider $\theta_{i,j} = p_i p_j - \lambda_{i,j}$. From Eq. (2.13), $\lambda_{i,j} = \lambda p_i = 3\lambda$, with the condition $(3\lambda)\mathbf{mod}p_j = (2n)\mathbf{mod}5$. Because $(2n)\mathbf{mod}5 \neq 0$, so $\lambda \leq 4$ and $\theta_{i,j} = p_i p_j - 3\lambda = 3(5 - \lambda) \geq 3$.
  $\left\lfloor \frac{ap_j+b+\theta_{i,j}}{3p_j} \right\rfloor \geq \left\lfloor \frac{2 \cdot 5+2+3}{3 \cdot 5} \right\rfloor = 1$.
  So, $\left\lfloor \frac{ap_j+b+\theta_{i,j}}{3p_j} \right\rfloor - \varepsilon_4 \geq 0 \quad \text{if} \quad s = 1$,
  or $\varepsilon' = s + \left\lfloor \frac{ap_j+b+\theta_{i,j}}{3p_j} \right\rfloor - \varepsilon_4 \geq 1$.

□

Therefore,

$$\begin{aligned}
\varepsilon &= \varepsilon' - a + \varepsilon'' + \left\lfloor \tfrac{ap_j+b+\theta_{j,i}}{3p_j} \right\rfloor + \left\lfloor \tfrac{ap_j+b+\theta_{ij}}{3p_j} \right\rfloor \\
&\geq 1 - a + \left\lfloor \tfrac{ap_j+b+\theta_{j,i}}{3p_j} \right\rfloor.
\end{aligned}$$

If $a \leq 1$ then $\varepsilon \geq 1 - a \geq 0$.
Else for $a = 2(< p_i)$, because $\theta_{j,i} = p_i p_j - \lambda p_j \geq p_j$ $(1 \leq \lambda \leq p_i - 1)$, $\left\lfloor \frac{ap_j+b+\theta_{j,i}}{p_i p_j} \right\rfloor \geq \left\lfloor \frac{2p_j+p_j}{3p_j} \right\rfloor = 1$. So $\varepsilon \geq 1 - a + 1 \geq 0$.

In summary, $\varepsilon \geq 0$ for all $p_i < p_j \leq \sqrt{m}$. the Lemma is proved. □

If $(2n)\mathbf{mod}p_i = 0$,

$$\begin{aligned}
\varepsilon &= m\left[1-\frac{1}{p_i}\right]\left[1-\frac{1}{p_j}-\frac{\widetilde{1}}{p_j}\right]-\left\lceil m\left(1-\frac{3}{p_j}\right)\right\rceil\left[1-\frac{1}{p_i}\right]\\
&= sp_ip_j\left(1-\frac{1}{p_i}\right)\left(1-\frac{2}{p_j}\right)-sp_ip_j\left(1-\frac{1}{p_i}\right)\left(1-\frac{3}{p_j}\right)\\
&\quad +t-\left\lfloor\frac{t}{p_i}\right\rfloor-\left\lfloor\frac{t}{p_j}\right\rfloor+\left\lfloor\frac{t}{p_ip_j}\right\rfloor-\left\lfloor\frac{t+\theta_j}{p_j}\right\rfloor+\left\lfloor\frac{t+\theta_{i,j}}{p_ip_j}\right\rfloor\\
&\quad -\left(t-\left\lfloor\frac{3t}{p_j}\right\rfloor\right)+\left\lfloor\frac{t-\left\lfloor\frac{3t}{p_j}\right\rfloor}{p_i}\right\rfloor\\
&= s(p_i-1)+\left\lfloor\frac{3t}{p_j}\right\rfloor-\left\lfloor\frac{t}{p_j}\right\rfloor-\left\lfloor\frac{t+\theta_j}{p_j}\right\rfloor+\left\lfloor\frac{t}{p_ip_j}\right\rfloor+\left\lfloor\frac{t+\theta_{i,j}}{p_ip_j}\right\rfloor-\left\lfloor\frac{\left\lfloor\frac{3t}{p_j}\right\rfloor}{p_i}\right\rfloor-\varepsilon_1\\
&= s(p_i-1)+3a+\left\lfloor\frac{3b}{p_j}\right\rfloor-a-a-\left\lfloor\frac{b+\theta_j}{p_j}\right\rfloor+\left\lfloor\frac{ap_j+b+\theta_{i,j}}{p_ip_j}\right\rfloor-\left\lfloor\frac{3a+\left\lfloor\frac{3b}{p_j}\right\rfloor}{p_i}\right\rfloor-\varepsilon_1\\
&= s(p_i-1)+a+\left\lfloor\frac{3b}{p_j}\right\rfloor+\left\lfloor\frac{ap_j+b+\theta_{i,j}}{p_ip_j}\right\rfloor-\varepsilon_1-\varepsilon_3-\varepsilon_4,
\end{aligned}$$

where

$$\begin{cases}
\varepsilon_1 &= \left\lfloor\frac{\left\lfloor\frac{3t}{p_j}\right\rfloor\mathbf{mod}p_i+\left(t-\left\lfloor\frac{3t}{p_j}\right\rfloor\right)\mathbf{mod}p_i}{p_i}\right\rfloor\le 1\\
\varepsilon_3 &= \left\lfloor\frac{3a+\left\lfloor\frac{3b}{p_j}\right\rfloor}{p_i}\right\rfloor\le 2\\
\varepsilon_4 &= \left\lfloor\frac{b+\theta_j}{p_j}\right\rfloor\le 1,
\end{cases}$$

$$\Delta\varepsilon=\varepsilon_1+\varepsilon_3+\varepsilon_4\le 4.$$

We can follow the same method above (it is a little easier),

$$\begin{cases}
m\left[1-\frac{1}{p_i}\right]\left[1-\frac{1}{p_j}-\frac{\widetilde{1}}{p_j}\right]\ge\left\lceil m\left(1-\frac{3}{p_j}\right)\right\rceil\left[1-\frac{1}{p_i}\right]\\
m\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right]\left[1-\frac{1}{p_j}\right]\ge\left\lceil m\left(1-\frac{3}{p_j}\right)\right\rceil\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right].
\end{cases}\tag{4.3}$$

Eq. (4.1) and (4.2) mean that, after deleted the items of $Z_k\mathbf{mod}p_j=0,\lambda_j$ from Z, the items of $Z_k\mathbf{mod}p_i\neq 0,\lambda_i$, will have,

$$S(m,p_i\nparallel 0,\lambda_i;p_j\nparallel 0,\lambda_j)=\quad S(\lceil m(1-3/p_j)\rceil,p_i\nparallel 0,\lambda_i)+\Delta(p_i\nparallel 0,\lambda_i),\tag{4.4}$$

where $\Delta(p_i\nparallel 0,\lambda_i)\ge 0$ is the extra items of $Z_k\mathbf{mod}p_i\neq 0,\lambda_i$ and $Z_k\mathbf{mod}p_j\neq 0,\lambda_j$. Thus the effect of $\left[1-\frac{1}{p_j}-\frac{\widetilde{1}}{p_j}\right]$ is that it constructs a new effective nature sequence with at least $\lceil m(1-3p_j)\rceil$ items which satisfy the condition $Z_k\mathbf{mod}p_j\neq 0,\lambda_j$.

This lemma means that, from equation (4.1) and (3.6), we can let $m\left[1-\frac{1}{p_j}-\frac{\widetilde{1}}{p_j}\right]=\left\lceil m\left(1-\frac{3}{p_j}\right)\right\rceil+t,t\ge 0$, and operated by $\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right]$ to attain the items which have no multiples of p_i and p_j.

$$\begin{aligned}
&m\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right]\left[1-\frac{1}{p_j}-\frac{\widetilde{1}}{p_j}\right]=S(m,p_i\nparallel 0,\lambda_i;p_j\nparallel 0,\lambda_j)\\
&=\left(\left\lceil m\left(1-\frac{3}{p_j}\right)\right\rceil+t\right)\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right]\\
&=S(\lceil m(1-3/p_j)\rceil,p_i\nparallel 0,\lambda_i)+\Delta(p_i\nparallel 0,\lambda_i)\\
&=\left\lceil m\left(1-\frac{3}{p_j}\right)\right\rceil\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right]+t\left[1-\frac{1'}{p_i}-\frac{1''}{p_i}\right]\\
&\ge\left\lceil m\left(1-\frac{3}{p_j}\right)\right\rceil\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right].
\end{aligned}\tag{4.5}$$

Lemma 4.3.

$$m\left[1-\tfrac{1}{2}\right]\left[1-\tfrac{1}{3}-\tfrac{\tilde{1}}{3}\right] \geq \left\lceil \tfrac{m}{6}\right\rceil - 1. \tag{4.6}$$

Proof. For $p_i = 2, p_j = 3$, $m = 6s + t$, from Eq. (3.4),

$$\begin{aligned}&m\left[1-\tfrac{1}{2}\right]\left[1-\tfrac{1}{3}-\tfrac{\tilde{1}}{3}\right] = 6s\left[1-\tfrac{1}{2}\right]\left[1-\tfrac{1}{3}-\tfrac{\tilde{1}}{3}\right] + t\left[1-\tfrac{1}{2}\right]\left[1-\tfrac{1}{3}-\tfrac{\tilde{1}}{3}\right]\\ &\geq 6s\left(1-\tfrac{1}{2}\right)\left(1-\tfrac{2}{3}\right) = s = \left\lfloor \tfrac{m}{6}\right\rfloor \geq \left\lceil \tfrac{m}{6}\right\rceil - 1.\end{aligned}$$

For the residual class of modulo 6, $X = \{6s+1, 6s+2, \cdots, 6s+6\}$, $X_k\mathbf{mod}6 = \{1, 2, 3, 4, 5, 6\}$, there are 4 elements $(2, 3, 4, 6)$ of multiples of 2 or 3. For the other elements $X_k\mathbf{mod}6 = \{1, 5\}$, there is at least one item with $3 \nmid (2n - X_k)$ or $X_k\mathbf{mod}3 \neq (2n)\mathbf{mod}3$. So m will have at least $\left\lfloor \frac{m}{6}\right\rfloor \geq \left\lceil \frac{m}{6}\right\rceil - 1$ double prime pairs. □

Lemma 4.4. *For* $i = 1, 2, \cdots, i_m, j$,

$$m\prod_{i=1}^{i_m}\left[1-\tfrac{1}{p_i}-\tfrac{\tilde{1}}{p_i}\right]\left[1-\tfrac{1}{p_j}-\tfrac{\tilde{1}}{p_j}\right] \geq \left\lceil m\left(1-\tfrac{3}{p_j}\right)\right\rceil \textstyle\prod_{i=1}^{i_m}\left[1-\tfrac{1}{p_i}-\tfrac{\tilde{1}}{p_i}\right] \tag{4.7}$$

Proof. From equation (2.20), (4.1), (4.2), (4.4), (4.5),

For $i \neq k$, we have $S(m, p_i \nparallel 0, \lambda_i; p_j \nparallel 0, \lambda_j) \geq S(\lceil m(1-3/p_j)\rceil, p_i \nparallel 0, \lambda_i)$

and $S(m, p_k \nparallel 0, \lambda_k; p_j \nparallel 0, \lambda_j) \geq S(\lceil m(1-3/p_j)\rceil, p_k \nparallel 0, \lambda_k)$,

so that $S(m, p_i \nparallel 0, \lambda_i; p_k \nparallel 0, \lambda_k; p_j \nparallel 0, \lambda_j) = S(\lceil m(1-3/p_j)\rceil, p_i \nparallel 0, \lambda_i; p_k \nparallel 0, \lambda_k) + \Delta(p_i \nparallel 0, \lambda_i; p_k \nparallel 0, \lambda_k) \geq S(\lceil m(1-3/p_j)\rceil, p_i \nparallel 0, \lambda_i; p_k \nparallel 0, \lambda_k)$.

Thus $S(m, p_{i_1} \nparallel 0, \lambda_{i_1}; p_{i_2} \nparallel 0, \lambda_{i_2}; p_j \nparallel 0, \lambda_j) \geq S(\lceil m(1-3/p_j)\rceil, p_{i_1} \nparallel 0, \lambda_{i_1}; p_{i_2} \nparallel 0, \lambda_{i_2})$. $\cdots\cdots$. $S(m, p_{i_1} \nparallel 0, \lambda_{i_1}; p_{i_2} \nparallel 0, \lambda_{i_2}; \cdots, p_{i_m} \nparallel 0, \lambda_{i_m}; p_j \nparallel 0, \lambda_j) \geq S(\lceil m(1-3/p_j)\rceil, p_{i_1} \nparallel 0, \lambda_{i_1}; p_{i_2} \nparallel 0, \lambda_{i_2}; \cdots, p_{i_m} \nparallel 0, \lambda_{i_m})$,

$$\begin{aligned}&m\prod_{i=1}^{i_m}\left[1-\tfrac{1}{p_i}-\tfrac{\tilde{1}}{p_i}\right]\left[1-\tfrac{1}{p_j}-\tfrac{\tilde{1}}{p_j}\right]\\ &= S(m, p_1 \nparallel 0, \lambda_1; p_2 \nparallel 0, \lambda_2; \cdots; p_{i_m} \nparallel 0, \lambda_{i_m}; p_j \nparallel 0, \lambda_j)\\ &\geq S(\lceil m(1-3/p_j)\rceil, p_1 \nparallel 0, \lambda_1; p_2 \nparallel 0, \lambda_2; \cdots; p_{i_m} \nparallel 0, \lambda_{i_m})\\ &= \left\lceil m\left(1-\tfrac{3}{p_j}\right)\right\rceil \prod_{i=1}^{i_m}\left[1-\tfrac{1}{p_i}-\tfrac{\tilde{1}}{p_i}\right].\end{aligned}$$

Suppose that for $1 < r \leq i_m$,

$$m\prod_{i=r}^{i_m}\left[1-\tfrac{1}{p_i}-\tfrac{\tilde{1}}{p_i}\right]\left[1-\tfrac{1}{p_j}-\tfrac{\tilde{1}}{p_j}\right] = \left\lceil m\left(1-\tfrac{3}{p_j}\right)\right\rceil \prod_{i=r}^{i_m}\left[1-\tfrac{1}{p_i}-\tfrac{\tilde{1}}{p_i}\right] + t, \tag{4.8}$$

where $t \geq 0$. It means that the effect of operator $\left[1-\frac{1}{p_j}-\frac{\tilde{1}}{p_j}\right]$ when operating on m is that dividing $Z = \{1, 2, \cdots, m\}$ into two sets $X = \{1, 2, \cdots, m' = \left\lceil m\left(1-\frac{3}{p_j}\right)\right\rceil\}$ and $X' = \{X'_1, X'_2, \cdots, X'_t, t = m - m' \geq 0\}$. From equation (2.22), (2.23), (4.1), (4.2), and (4.8), we have

$$\begin{aligned}
& m\prod_{i=r-1}^{i_m}\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right]\left[1-\tfrac{1}{p_j}-\tfrac{\widetilde{1}}{p_j}\right] \\
&= m\prod_{i=r}^{i_m}\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right]\left[1-\tfrac{1}{p_j}-\tfrac{\widetilde{1}}{p_j}\right]\left[1-\tfrac{1}{p_{r-1}}-\tfrac{\widetilde{1}}{p_{r-1}}\right] \\
&= \left(\left\lceil m\left(1-\tfrac{3}{p_j}\right)\right\rceil\prod_{i=r}^{i_m}\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right]+t\right)\left[1-\tfrac{1'}{p_{r-1}}-\tfrac{1''}{p_{r-1}}\right] \\
&\geq \left\lceil m\left(1-\tfrac{3}{p_j}\right)\right\rceil\prod_{i=r}^{i_m}\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right]\left[1-\tfrac{1'}{p_{r-1}}-\tfrac{1''}{p_{r-1}}\right] \\
&= \left\lceil m\left(1-\tfrac{3}{p_j}\right)\right\rceil\prod_{i=r-1}^{i_m}\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right].
\end{aligned}$$

or

$$\begin{aligned}
& m\prod_{i=r}^{i_m}\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right]\left[1-\tfrac{1}{p_j}-\tfrac{\widetilde{1}}{p_j}\right]\left[1-\tfrac{1}{p_{r-1}}-\tfrac{\widetilde{1}}{p_{r-1}}\right] \\
&= m\prod_{i=r}^{i_m}\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right]\left[1-\tfrac{1}{p_j}-\tfrac{\widetilde{1}}{p_j}\right]-m\prod_{i=r}^{i_m}\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right]\left[1-\tfrac{1}{p_j}-\tfrac{\widetilde{1}}{p_j}\right]\left[\tfrac{1}{p_{r-1}}\right] \\
&\quad -m\prod_{i=r}^{i_m}\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right]\left[1-\tfrac{1}{p_j}-\tfrac{\widetilde{1}}{p_j}\right]\left[\tfrac{\widetilde{1}}{p_{r-1}}\right] \\
&= \left\lceil m\left(1-\tfrac{3}{p_j}\right)\right\rceil\prod_{i=r}^{i_m}\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right]+t \\
&\quad -\left\lceil m\left(1-\tfrac{3}{p_j}\right)\right\rceil\prod_{i=r}^{i_m}\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right]\left[\tfrac{1}{p_{r-1}}\right]-t\left[\tfrac{1'}{p_{r-1}}\right] \\
&\quad -\left\lceil m\left(1-\tfrac{3}{p_j}\right)\right\rceil\prod_{i=r}^{i_m}\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right]\left[\tfrac{\widetilde{1}}{p_{r-1}}\right]-t\left[\tfrac{1''}{p_{r-1}}\right] \\
&= \left\lceil m\left(1-\tfrac{3}{p_j}\right)\right\rceil\prod_{i=r}^{i_m}\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right]\left[1-\tfrac{1}{p_{r-1}}-\tfrac{\widetilde{1}}{p_{r-1}}\right]+t\left[1-\tfrac{1'}{p_{r-1}}-\tfrac{1''}{p_{r-1}}\right] \\
&\geq \left\lceil m\left(1-\tfrac{3}{p_j}\right)\right\rceil\prod_{i=r-1}^{i_m}\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right].
\end{aligned}$$

If $(2n)\mathbf{mod}p_{r-1}=0$, then

$$\begin{aligned}
& m\prod_{i=r-1}^{i_m}\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right]\left[1-\tfrac{1}{p_j}-\tfrac{\widetilde{1}}{p_j}\right] \\
&\geq \left\lceil m\left(1-\tfrac{3}{p_j}\right)\right\rceil\prod_j\left[1-\tfrac{1}{p_{r-1}}\right]\prod_{i=r}^{i_m}\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right].
\end{aligned}\tag{4.9}$$

□

In fact, if

$$m\prod_{i=1}^{i_m}\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right]\left[1-\tfrac{1}{p_j}-\tfrac{\widetilde{1}}{p_j}\right]<\left\lceil m\left(1-\tfrac{3}{p_j}\right)\right\rceil\textstyle\prod_{i=1}^{i_m}\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right],$$

then for any p_i, p_j, before deleting the multiples of other primes, it must have

$$m\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right]\left[1-\tfrac{1}{p_j}-\tfrac{\widetilde{1}}{p_j}\right]<\left\lceil m\left(1-\tfrac{3}{p_j}\right)\right\rceil\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right].$$

which contradicts Lemma 4.2. So this lemma is true. □

With Lemma 4.2, the operator $\left[1-\frac{1}{p_j}-\frac{\widetilde{1}}{p_j}\right]$ can be represented by $\left(1-\frac{3}{p_j}\right)$, and other operator $\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right]$ can operate on this inequality unchanged.

5. EXPLANATION

Let $m = sp_ip_j + ap_j + b \geq p_j^2$, then for all $p_i < p_j$, the effect of $m\left[1-\frac{1}{p_j}-\frac{\widetilde{1}}{p_j}\right]$ is a nature sequence $X = \{f(Z)\}$ whose number is not less than $\lceil m(1-3/p_j)\rceil$. The reason is as follows. When a nature sequence is deleted by the multiples of p_j, the sequence is subtracted by $\left[\frac{m}{p_j}\right]+\left[\frac{m+\theta_j}{p_j}\right]$. We can arrange the m items in a table of p_j rows (Table 1). $\left[\frac{m}{p_j}\right]$ will delete the p_jth row, and $\left[\frac{m+\theta_j}{p_j}\right]$ will delete the $(2n)\mathbf{mod}p_jth$ row. Thus there are (p_j-2) rows left in which each item $Z_k\mathbf{mod}p_j \neq 0, (2n)\mathbf{mod}p_j$.

TABLE 1. Set Z

1	p_j+1	$\cdots$	$(p_i-1)p_j+1$	$\cdots$	$(sp_i-1)p_j+1$	sp_ip_j+1	$\cdots$	$sp_ip_j+ap_j+1$
2	p_j+2	$\cdots$	$(p_i-1)p_j+2$	$\cdots$	$(sp_i-1)p_j+2$	sp_ip_j+2	$\cdots$	$sp_ip_j+ap_j+2$
$\cdots$	$\cdots$	$\cdots$	$\cdots$	$\cdots$	$\cdots$	$\cdots$	$\cdots$	$\cdots$
b	p_j+b	$\cdots$	$(p_i-1)p_j+b$	$\cdots$	$(sp_i-1)p_j+b$	sp_ip_j+b	$\cdots$	$sp_ip_j+ap_j+b$
$\cdots$	$\cdots$	$\cdots$	$\cdots$	$\cdots$	$\cdots$	$\cdots$	$\cdots$	
p_j	$2p_j$	$\cdots$	p_ip_j	$\cdots$	sp_ip_j	$sp_ip_j+p_j$	$\cdots$	

But every p_i items $(0 \leq r = Z_k\mathbf{mod}p_i \leq p_i-1)$ in any row of the first sp_i columns consist in a complete system of residues modulo p_i, because $C_1 = \{1, p_j+1, 2p_j+1, \cdots, (p_i-1)p_j+1\}$ and $C_r = \{C_1+r\}$ are both complete system of residues modulo p_i, where r is any (row or column) constant. There are (p_j-2) such rows or $sp_i(p_j-2)$ items left. These items are effective to a nature sequence when deleting multiples of p_i,

$$sp_ip_j\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right]\left[1-\frac{1}{p_j}-\frac{\widetilde{1}}{p_j}\right] = sp_i(p_j-2)\left[1-\frac{1}{p_i}-\frac{\widetilde{1}}{p_i}\right] = s(p_i-2)(p_j-2).$$

Let $t = ap_j+b$, $0 \leq b \leq p_j-1$, $0 \leq a \leq p_i-1$, $t\left[\frac{1}{p_j}+\frac{\widetilde{1}}{p_j}\right]$ will delete at most $a+(a+1) \leq t\frac{1}{p_j}+sp_ip_j\frac{1}{p_j} = \frac{m}{p_j}$ items. If we add these items by removing those from the end of sequence, then the sequence is again effective to a nature sequence, which has at least,

$$\begin{aligned}
&M(j) \geq sp_i(p_j-2) + t\left[1-\frac{1}{p_j}-\frac{\widetilde{1}}{p_j}\right] - [a+(a+1)] \\
&\geq sp_i(p_j-3) + t + sp_i - 4a - 2 \\
&= sp_i(p_j-3) - 3\left\lfloor\frac{t}{p_j}\right\rfloor + t + (sp_i-a-2) \\
&\geq sp_ip_j(1-\frac{3}{p_j}) + \left\lceil t-\frac{3t}{p_j}\right\rceil + (sp_i-a-2) \\
&= \left\lceil (sp_ip_j+t)(1-\frac{3}{p_j})\right\rceil + (sp_i-a-2) \\
&= \left\lceil m(1-\frac{3}{p_j})\right\rceil + (sp_i-a-2).
\end{aligned}$$

For $s \geq 2$ or $a \leq p_i - 2$, we have $(sp_i - a - 2) = (s-1)p_i + (p_i - a - 1) - 1 \geq 0$, $M(j) \geq \left\lceil m(1 - \frac{3}{p_j}) \right\rceil$.

For $s = 1$ and $a = p_i - 1$, the items of t have p_j rows, $p_i - 1$ columns and some b items. In each of the first b rows, there are exact p_i items which consist in a complete system of residues modulo p_i, and these items can be considered as an effective nature sequence when deleting the multiples of p_i $(Z_k\mathbf{mod}p_i = 0, \lambda_i)$. The other items have at most p_j rows and $p_i - 1$ columns where the multiples of p_j have at most $2(p_i - 1)$. As before, we can add these items to make the t as an effective nature sequence, therefore,

$M(j) \geq sp_i(p_j - 2) + t\left[1 - \frac{1}{p_j} - \frac{\tilde{1}}{p_j}\right] - 2(p_i - 1) \geq \left\lceil m(1 - \frac{3}{p_j}) \right\rceil + (sp_i - a - 1) \geq \left\lceil m(1 - \frac{3}{p_j}) \right\rceil$.

Thus for any $p_i < p_j$, the original sequence of $m \geq p_j^2$, after deleted all the multiples of p_j from $2n = Z_k + Z_k'$, is effective to reconstruct a new nature sequence having at least $\left\lceil m(1 - \frac{3}{p_j}) \right\rceil$ items.

Example 5.1. $2n = 46$, $m = 2n - 1 = 45$, $Z = \{1, 2, \cdots, 45\}$,

For $p_j = 5$, after deleted the item of $Z_k\mathbf{mod}p_j = 0, (2n)\mathbf{mod}p_j$, it becomes $Z \to Z' = \{2, 3, 4, 7, 8, 9, 12, 13, 14, 17, 18, 19, 22, 23, 24, 27, 28, 29, 32, 33, 34, 37, 38,$-$39, 42, 43, 44\}$. we can rearrange these items as $Z' = \{(19), 2, 3, 4, (23)7, 8, 9, (22),$-$(29), 12, 13, 14, (27), (28), 17, 18, ||, 24, 32, 33, 34, 37, 38, 39, 42, 43, 44\}$. The first $\left\lceil m\left(1 - \frac{3}{p_j}\right) \right\rceil = 18$ items can be taken as an effective nature sequence $(\{1, 2, \cdots, 18\})$ from the original one when deleting the items of $Z_k\mathbf{mod}3 = 0, (2n)\mathbf{mod}3$. The other sequence $X = \{24, 32, 33, 34, 37, 38, 39, 42, 43, 44\}$, having at least zero item after deleted the multiples of all primes, will be neglected in further process.

6. Proof of Theorem 1.1

For a given 2n, consider the possible pairs of $2n = p_i + p_j$, where p_i and p_j are both primes.

Lemma 6.1. *The (double) prime pairs in* $2n$*,*

$$D(2n) \geq \left\lceil \frac{p_v}{6} \prod_{i=3}^{v-1} \frac{p_{i+1} - 3}{p_i} - \frac{3}{2} \right\rceil = \left\lceil \frac{W(v) - 9}{6} \right\rceil, \tag{6.1}$$

where

$$W(v) = p_v \prod_{i=3}^{v-1} \frac{p_{i+1} - 3}{p_i}. \tag{6.2}$$

If $W(v) > 9$ *then* $D(2n) \geq 1$.

Proof. From equation (2.9), (3.3), (4.7), (4.9) and (4.6),

$$
\begin{aligned}
D_0(m) &= m\prod_{(2n)\ \textbf{mod}\ p_i=0}\left[1-\tfrac{1}{p_i}\right]\prod_{(2n)\ \textbf{mod}\ p_j\neq 0}\left[1-\tfrac{1}{p_j}-\tfrac{\widetilde{1}}{p_j}\right]\\
&\geq m\left[1-\tfrac{1}{2}\right]\prod_{i=2}^{v}\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right]\\
&\geq \left\lceil m\left(1-\tfrac{3}{p_v}\right)\right\rceil\left[1-\tfrac{1}{2}\right]\prod_{i=2}^{v-1}\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right]\\
&\geq \left\lceil\left\lceil m\left(1-\tfrac{3}{p_v}\right)\right\rceil\left(1-\tfrac{3}{p_{v-1}}\right)\right\rceil\left[1-\tfrac{1}{2}\right]\prod_{i=2}^{v-2}\left[1-\tfrac{1}{p_i}-\tfrac{\widetilde{1}}{p_i}\right]\\
&\geq \cdots\\
&\geq \left\lceil m\prod_{i=3}^{v}\left(1-\tfrac{3}{p_i}\right)\right\rceil\left[1-\tfrac{1}{2}\right]\left[1-\tfrac{1}{3}-\tfrac{\widetilde{1}}{3}\right]\\
&\geq \left\lceil \tfrac{m}{6}\prod_{i=3}^{v}\left(1-\tfrac{3}{p_i}\right)\right\rceil-1,
\end{aligned}
$$

but

$$
\begin{aligned}
&\prod_{i=3}^{v}\left(1-\tfrac{3}{p_i}\right)=\tfrac{p_3-3}{p_3}\tfrac{p_4-3}{p_4}\cdots\tfrac{p_{v-1}-3}{p_{v-1}}\tfrac{p_v-3}{p_v}\\
&=\tfrac{p_3-3}{p_v}\tfrac{p_4-3}{p_3}\tfrac{p_5-3}{p_4}\cdots\tfrac{p_v-3}{p_{v-1}}=\tfrac{2}{p_v}\prod_{i=3}^{v-1}\tfrac{p_{i+1}-3}{p_i},
\end{aligned}
$$

and $m=2n-1\geq p_v^2$, so

$$
D_0(2n-1)\geq\left\lceil\frac{p_v^2}{6}\frac{2}{p_v}\prod_{i=3}^{v-1}\frac{p_{i+1}-3}{p_i}\right\rceil-1=\left\lceil\frac{p_v}{3}\prod_{i=3}^{v-1}\frac{p_{i+1}-3}{p_i}\right\rceil-1 \tag{6.3}
$$

From equation (2.14), (2.15),

$$
\begin{aligned}
D(2n) &= \left\lceil\tfrac{D_0(2n-1)}{2}\right\rceil+D(\sqrt{2n})-D_1\geq\left\lceil\tfrac{1}{2}\left\lceil\tfrac{p_v}{3}\prod_{i=3}^{v-1}\tfrac{p_{i+1}-3}{p_i}\right\rceil-\tfrac{1}{2}\right\rceil-1\\
&\geq\left\lceil\tfrac{p_v}{6}\prod_{i=3}^{v-1}\tfrac{p_{i+1}-3}{p_i}-\tfrac{3}{2}\right\rceil.
\end{aligned}
$$

□

Proof of Theorem 1.1. Because for $v=5, p_v=11$, $W(v)=11\frac{4}{5}\frac{8}{7}=10.057>9$.
Suppose that for v, $W(v)=p_v\prod_{i=3}^{v-1}\frac{p_{i+1}-3}{p_i}>9$, then for $v+1$,

$$
\begin{aligned}
W(v+1) &= p_{v+1}\prod_{i=3}^{v}\tfrac{p_{i+1}-3}{p_i}=p_v\prod_{i=3}^{v-1}\tfrac{p_{i+1}-3}{p_i}\tfrac{p_{v+1}}{p_v}\tfrac{p_{v+1}-3}{p_v}\\
&= W(v)\tfrac{p_{v+1}^2-3p_{v+1}}{p_v^2}.
\end{aligned}
\tag{6.4}
$$

Because, $p_{v+1}=p_v+2\Delta, \Delta\geq 1, p_v\geq 11$,

$$
\begin{aligned}
&p_{v+1}^2-3p_{v+1}-p_v^2=(p_v+2\Delta)^2-3(p_v+2\Delta)-p_v^2\\
&=p_v^2+4\Delta p_v+4\Delta^2-3p_v-6\Delta-p_v^2\\
&=3(\Delta-1)p_v+\Delta(p_v+4\Delta-6)>0.
\end{aligned}
$$

Therefore, $\frac{p_{v+1}^2-3p_{v+1}}{p_v^2}>1$, $W(v+1)>W(v)>9$. From 'the principle of mathematical induction', we can conclude that for any $v\geq 5$, we have $W(v)\geq 9$ and $D(2n)\geq 1$, or there is at least one pair of double primes p_i, p_j such that $2n=p_i+p_j$ for any $2n>p_v^2+1=11^2+1=122$.

Besides, when $6 \leq 2n \leq 120$, we know that there is at least one pair of primes p_i, p_j such that $2n = p_i + p_j$. In fact, from Eq. (6.1), $D(2n)$ approaches infinity as n grows without bound. The proof is completed. □

Corollary 6.2. *Any odd number not less than 9 can be expressed as the sum of three odd primes.*

Proof. If n is odd number, $p_1 = 3$, then $n - p_1 \geq 6$ and can be represent as the sum of two primes $p_2 + p_3$ from Theorem 1.1. So $n = p_1 + p_2 + p_3$. □

Example 6.3 (Actual vs. Simplified Formula)**.** Figure 1 shows the minimum actual prime pairs $D(2n)$ (solid line) of $2n$ in the range of $[p_v^2 + 1, p_{v+1}^2 - 1]$ and the simplified formula (dashed line) from Eq. (6.1) against v. From this figure, it is easily seen that

$$D(2n) \geq \left\lceil \frac{p_v}{6} \prod_{i=3}^{v-1} \frac{p_{i+1}-3}{p_i} - \frac{3}{2} \right\rceil \geq 1. \tag{6.5}$$



FIGURE 1. The minimum number of actual prime pairs (solid) and its simplified formula (dashed) against v.

REFERENCES

[1] J. R. Chen., On the Representation of a Large Even Integer as the Sum of a Prime and the Product of at Most Two Primes., Sci. Sinica 16 (1973), 157-176.
[2] J. R. Chen., On the Representation of a Large Even Integer as the Sum of a Prime and the Product of at Most Two Primes. II., Sci. Sinica 16 (1978), 421-430.
[3] J. R. Chen & C. D. Pan., The exceptional set of Goldbach numbers, Sci. Sinica, 23 (1980), 416-430.
[4] H. L. Montgomery & R. C. Vaughan., The exceptional set in Goldbach's probem, Acta Arith, 27 (1975), 353-370.
[5] Kenneth H. Rosen, Elementanry Number Theory and Its Applications, Fifth Edition, Pearson Education Asia Limited and China Maxhine Press, 2005.

CHINESE ACADEMY OF SCIENCES, 99 DONGGANG RD. SHIJIAZHUANG, HEBEI, 050031, CHINA
E-mail address: `shouyudu@yahoo.com.cn`

CHINESE ACADEMY OF SCIENCES, A20 DATUN RD. CHAOYANG DST. BEIJING 100012, CHINA
E-mail address: `zldu@bao.ac.cn`